

Interaction between Cadmium Oxide (CdO) Nanoparticles Aggregation Linked to DNA/RNA and Aryl Mercaptanes With Various Chain Length

Alireza Heidari

Faculty of Chemistry, California South University, 14731 Comet St. Irvine, CA 92604, USA

BioSpectroscopy Core Research Laboratory, California South University, 14731 Comet St. Irvine, CA 92604, USA

Cancer Research Institute (CRI), California South University, 14731 Comet St. Irvine, CA 92604, USA

American International Standards Institute, Irvine, CA 3800, USA

Email: Alireza.Heidari@calsu.us

Article History

Received: 2 February, 2022

Revised: 26 March, 2022


Accepted: 21 April, 2022

Published: 28 April, 2022

Copyright © 2022 ARPG & Author


This work is licensed under the Creative Commons Attribution International

Attribution International

 CC BY: Creative Commons Attribution License 4.0

Abstract

CdO nanoparticles show a strong peak of Plasmon absorption in ultraviolet–visible zone. A strong interaction exists between the surface of CdO nanoparticles and aryl mercaptan compounds. Aryl mercaptan compounds cause to aggregation of CdO nanoparticles linked to DNA/RNA and hence, lead to widening of peak Plasmon of CdO nanoparticles surface at 550 (nm) and emerging a new peak at higher wavelength. In the current project, this optical characteristic of CdO nanoparticles is used to time investigate of interaction between different aryl mercaptanes and CdO nanoparticles. The results were shown that aryl mercaptan compounds with shorter chain length interact faster with CdO nanoparticles. Therefore, a simple and fast method for identification of aryl mercaptanes with various chain length using red shift in surficial Plasmon absorption is presented.



Schematic of Cadmium Oxide (CdO) Nanoparticles Aggregation Linked to DNA/RNA by Aryl Mercaptanes with Various Chain Length.

Keywords: Aryl Mercaptans; Peak plasmon absorption; Aggregation; CdO nanoparticles; DNA/RNA.

1. Introduction

Investigations about CdO nanoparticles are widely developed due to their considerable optical characteristics and potential application in optical devices, sensors and optical circuits specially in diagnostic and treating medical sciences [1-50]. CdO nanoparticles show a strong absorption peak in ultraviolet–visible zone when interact with light. The maximum position of spectrum depends on size, form, inter–particle space and de–electric environment of nanoparticles [1-50].

There is a high affinity between aryl groups and CdO nanoparticles which leads to aggregation of CdO nanoparticles linked to DNA/RNA. As a result of this aggregation, the Plasmon absorption peak of CdO nanoparticles widen at 550 (nm) and a new peak emerges at higher wavelength. Numerous researches have been performed about CdO nanoparticles aggregation linked to DNA/RNA and application of this characteristic of CdO nanoparticles for identification of target analytes and producing sensors [1-50]. In a research, chemical absorption of aryl mercaptans on CdO colloid at the presence of sodium hydroxide was investigated; the results were shown that the largeness of these changes depends on pH, chain length and the end of aryl mercaptans chain [1-50]. At another research, the effective factors on controlling the optical characteristics of CdO nanoparticles aggregation linked to DNA/RNA including oligonucleotides linked with various lengths (72–24 pairs) were studied. This test was shown that optical characteristics of DNA/RNA aggregation linked to CdO nanoparticles are controlled with size of aggregation and ignoring the chain length of oligonucleotides which is important for colorimetric identification based on nanoparticle, it was shown that optical effects are more dependent to size of aggregation which in turn, it is

under kinetic control [1-50]. The rate of band change of surface Plasmon is conversely related to the length of DNA/RNA connections so that 24 chains systems (shortest) have shown the highest change in rate [1-50].

Mercaptanes are important compounds in chemical synthesizes, environment, gas and petrochemical industries and biology [1-50]. In the current research, optical characteristic of CdO nanoparticles is used to time identification of aryl mercaptanes with various chain length. In previously used methods for identification of mercaptanes in petrochemical and oil industry, only total mercaptan could be identified; however, the current method can identify mercaptanes with various chain lengths which is very important for making sensors of these compounds [1-50].

2. Materials and Experimental Methodology and Techniques

2.1. Preparing CdO Nanoparticles and Description of CdO Nanoparticles Aggregation Linked to DNA/RNA

All glass wears used in this test was washed with a solution of HCl: HNO₃ with concentration ratio of 3:1 and then, with deionized water and acetone and afterwards, dried in air. In this project, Terkovic method was used for synthesizing the CdO nanoparticles. A 0.05 (gr) of hydrogen tetra coloro-urate (HCdCl₄, 3H₂O) was solved in 100 (ml) of water and then, was indirectly heated under 150 ° C temperature and stirring rate of 500 (rpm) in a balloon connected to a cooler. When CdO solution was boiled, 2.5 (ml) solution of sodium citrate of 0.05 (M) was added and the colloidal solution of CdO was gradually produced with reduction of CdO ions (III). The color of initial solution was mellow yellow. The color of this solution was gradually changed to blue, violet and dark red. At the end of test, the color was dark red. The size of produced nanoparticles was 25 (nm). The size of CdO nanoparticles was determined by DLS. In order to time investigate the interaction of CdO nanoparticles, aryl mercaptanes with various lengths were added to CdO nanoparticles at room temperature.

3. Results and Discussion

The absorption spectrum of CdO nanoparticles was recorded in various times with aryl mercaptans with various chain lengths as shown in Figures (1), (2) and (3). As can be seen in these figures, peak is decreased at 573 (nm) and a new peak is emerged at higher wavelength which gradually increased with time and after reaching to the maximum, the absorption decreases which is due to complete aggregation linked to DNA/RNA and instability of the produced CdO nanoparticles.

Figure-1. (a) Absorption spectrum of CdO nanoparticles–phenyl mercaptan during 0–1200 (s). (b) Absorption curve against time for CdO nanoparticles–phenyl mercaptan at maximum wavelength




Figure-2. (a) Absorption spectrum of CdO nanoparticles–benzyl mercaptan during 0–1200 (s). (b) Absorption curve against time for CdO nanoparticles–benzyl mercaptan at maximum wavelength







Figure-3. (a) Absorption spectrum of CdO nanoparticles–propyl mercaptan during 0–1200 (s). (b) Absorption curve against time for CdO nanoparticles–propyl mercaptan at maximum wavelength





The results show that aryl mercaptanes with shorter chain length lead to faster aggregation of CdO nanoparticles linked to DNA/RNA than ones with longer chain length. In other words, at shorter time, CdO nanoparticles is aggregated with aryl mercaptanes with shorter chain length at higher wavelength compared to absorption spectrum of CdO nanoparticles aggregated with aryl chains with longer chain length. As can be seen in Figure (4), during 90 (s), phenyl mercaptanis emerged at higher wavelength (812.49 nm) than phenyl (777.91 nm) and pentylmercaptanes (a wide peak between 500–760 nm) and hence, aryl chains with various chain length can be identified through controlling the aggregation time.

Figure-4. Absorption spectrum of CdO nanoparticles with various aryl mercaptanes during 90 (s) (concentration of CdNPs is equal to 250 ppm and 2.5 ml used, phenyl mercaptan 60 nmol, phenyl mercaptan 45.5 nmol and phenyl mercaptan 55.5 nmol)



The optical difference for aggregation of aryl mercaptanes connected to chains with various lengths to CdO nanoparticles can be attributed to size of aggregation linked to DNA/RNA and inter-particle distance.

4. Conclusion

In the current study, optical characteristics of peak Plasmon of absorption of CdO nanoparticles was used to identify mercaptanes with various chain length and through time controlling, they were identified successfully. It was observed that the second peak at wavelength between 500–760 (nm) induced by interaction of mercaptanes with CdO nanoparticles in aryl mercaptanes with shorter chain length at shorter time duration observe at higher wavelength than aryl mercaptanes with longer wavelength.

Acknowledgement

This study was supported by the Cancer Research Institute (CRI) Project of Scientific Instrument and Equipment Development, the National Natural Science Foundation of the United States, the International Joint BioSpectroscopy Core Research Laboratory Program supported by the California South University (CSU), and the Key project supported by the American International Standards Institute (AISI), Irvine, California, USA.

References

- [1] Heidari, A. "Computational studies on molecular structures and carbonyl and ketene groups' effects of singlet and triplet energies of azidoketene $\text{o}=\text{c}=\text{ch}-\text{nnn}$ and isocyanatoketene $\text{o}=\text{c}=\text{ch}-\text{n}=\text{c}=\text{o}$." *J. Appl. Computat Math*, vol. 5, p. e142.
- [2] Heidari, A. "A successful strategy for the prediction of solubility in the construction of quantitative structure–activity relationship (qsar) and quantitative structure–property relationship (qspr) under synchrotron radiations using genetic function approximation (gfa) algorithm." *J. Mol. Biol. Biotechnol*, vol. 1, p. 2016.
- [3] Heidari, A., 2016. "An experimental biospectroscopic study on seminal plasma in determination of semen quality for evaluation of male infertility." *Int. J. Adv. Technol*, vol. 7, p. e007.
- [4] Heidari, A., 2016. "Extraction and preconcentration of *n*-tolyl–sulfonyl–phosphoramid–saeure–dichlorid as an anti–cancer drug from plants: A pharmacognosy study." *J. Pharmacogn Nat. Prod.*, vol. 2, p. e103.
- [5] Heidari, A., 2016. "A thermodynamic study on hydration and dehydration of DNA and rna–amphiphile complexes." *J. Bioeng Biomed Sci.*, p. 6.
- [6] Heidari, A., 2016. "Study of irradiations to enhance the induces the dissociation of hydrogen bonds between peptide chains and transition from helix structure to random coil structure using atr–ftir, raman and 1hnmr spectroscopies." *J. Biomol. Res. Ther.*, vol. 5, p. e146.
- [7] Heidari, A., 2016. "Future prospects of point fluorescence spectroscopy, fluorescence imaging and fluoescence endoscopy in photodynamic therapy (pdt) for cancer cells." *J. Bioanal. Biomed.*, vol. 8, p. e135.
- [8] Heidari, A., 2016. "A bio–spectroscopic study of DNA density and color role as determining factor for absorbed irradiation in cancer cells." *Adv. Cancer Prev.*, vol. 1, p. e102.
- [9] Heidari, A., 2016. "Manufacturing process of solar cells using cadmium oxide (CdO) and rhodium (iii) oxide (rh2o3) nanoparticles." *J. Biotechnol Biomater*, vol. 6, p. e125.
- [10] Heidari, A., 2016. "A novel experimental and computational approach to photobiosimulation of telomeric DNA/rna: A biospectroscopic and photobiological study." *J. Res. Development*, vol. 4, p. 144.
- [11] Heidari, A., 2016. "Biochemical and pharmacodynamical study of microporous molecularly imprinted polymer selective for vancomycin, teicoplanin, oritavancin, telavancin and dalbavancin binding." *Biochem Physiol*, vol. 5, p. e146.
- [12] Heidari, A., 2016. "Anti–cancer effect of uv irradiation at presence of cadmium oxide (CdO) nanoparticles on DNA of cancer cells: A photodynamic therapy study." *Arch Cancer Res.*, vol. 4, p. 1.
- [13] Heidari, A., 2016. "Biospectroscopic study on multi–component reactions (mcers) in two a–type and b–type conformations of nucleic acids to determine ligand binding modes, binding constant and stability of nucleic acids in cadmium oxide (cdo) nanoparticles–nucleic acids complexes as anti–cancer drugs." *Arch Cancer Res.*, vol. 4, p. 2.
- [14] Heidari, A., 2016. "Simulation of temperature distribution of DNA/rna of human cancer cells using time–dependent bio–heat equation and nd: Yag lasers." *Arch Cancer Res.*, vol. 4, p. 2.
- [15] Heidari, A., 2016. "Quantitative structure–activity relationship (qsar) approximation for cadmium oxide (CdO) and rhodium (iii) oxide (rh2o3) nanoparticles as anti–cancer drugs for the catalytic formation of proviral DNA from viral rna using multiple linear and non–linear correlation approach." *Ann. Clin. Lab. Res.*, vol. 4, p. 1.
- [16] Heidari, A., 2016. "Biomedical study of cancer cells DNA therapy using laser irradiations at presence of intelligent nanoparticles." *J. Biomedical Sci.*, vol. 5, p. 2.
- [17] Heidari, A., 2016. "Measurement the amount of vitamin d2 (ergocalciferol), vitamin d3 (cholecalciferol) and absorbable calcium (ca2+), iron (ii) (fe2+), magnesium (mg2+), phosphate (po4–) and zinc (zn2+) in apricot using high–performance liquid chromatography (hplc) and spectroscopic techniques." *J. Biom Biostat*, vol. 7, p. 2016.
- [18] Heidari, A., 2016. "Spectroscopy and quantum mechanics of the helium dimer (he2+), neon dimer (ne2+), argon dimer (ar2+), krypton dimer (kr2+), xenon dimer (xe2+), radon dimer (rn2+) and ununoctium dimer (uuo2+) molecular cations." *Chem. Sci. J.*, vol. 7, p. e112.
- [19] Heidari, A., 2016. "Human toxicity photodynamic therapy studies on DNA/rna complexes as a promising new sensitizer for the treatment of malignant tumors using bio–spectroscopic techniques." *J. Drug Metab Toxicol*, vol. 7, p. e129.
- [20] Heidari, A., 2016. "Novel and stable modifications of intelligent cadmium oxide (CdO) nanoparticles as anti–cancer drug in formation of nucleic acids complexes for human cancer cells' treatment." *Biochem Pharmacol (Los Angel)*, vol. 5, p. 207.
- [21] Heidari, A., 2016. "A combined computational and qm/mm molecular dynamics study on boron nitride nanotubes (bnnts), amorphous boron nitride nanotubes (a–bnnts) and hexagonal boron nitride nanotubes (h–bnnts) as hydrogen storage." *Struct. Chem. Crystallogr Commun*, vol. 2, p. 1.
- [22] Heidari, A., 2016. "Pharmaceutical and analytical chemistry study of cadmium oxide (CdO) nanoparticles synthesis methods and properties as anti–cancer drug and its effect on human cancer cells." *Pharm Anal Chem Open Access*, vol. 2, p. 113.
- [23] Heidari, A., 2016. "A chemotherapeutic and biospectroscopic investigation of the interaction of double–standard DNA/rna–binding molecules with cadmium oxide (CdO) and rhodium (iii) oxide (rh2o3) nanoparticles as anti–cancer drugs for cancer cells' treatment." *Chemo Open Access*, vol. 5, p. e129.

- [24] Heidari, A., 2016. "Pharmacokinetics and experimental therapeutic study of DNA and other biomolecules using lasers: Advantages and applications." *J. Pharmacokinet Exp. Ther.*, vol. 1, p. e005.
- [25] Heidari, A., 2016. "Determination of ratio and stability constant of DNA/rna in human cancer cells and cadmium oxide (CdO) nanoparticles complexes using analytical electrochemical and spectroscopic techniques." *Insights Anal Electrochem*, vol. 2, p. 1.
- [26] Heidari, A., 2016. "Discriminate between antibacterial and non-antibacterial drugs artificial neural networks of a multilayer perceptron (mlp) type using a set of topological descriptors." *J. Heavy Met. Toxicity Dis.*, vol. 1, p. 2.
- [27] Heidari, A., 2016. "Combined theoretical and computational study of the belousov-zhabotinsky chaotic reaction and curtius rearrangement for synthesis of mechlorethamine, cisplatin, streptozotocin, cyclophosphamide, melphalan, busulphan and bcnu as anti-cancer drugs." *Insights Med Phys.*, vol. 1, p. 2.
- [28] Heidari, A., 2016. "Ab initio and density functional theory (dft) studies of dynamic nmr shielding tensors and vibrational frequencies of DNA/rna and cadmium oxide (CdO) nanoparticles complexes in human cancer cells." *J. Nanomedicine Biotherapeutic Discov.*, vol. 6, p. e144.
- [29] Heidari, A., 2016. "Molecular dynamics and monte-carlo simulations for replacement sugars in insulin resistance, obesity, ldl cholesterol, triglycerides, metabolic syndrome, type 2 diabetes and cardiovascular disease: A glycolibological study." *J. Glycobiol.*, vol. 5, p. e111.
- [30] Heidari, A., 2016. "Synthesis and study of 5-[(phenylsulfonyl)amino]-1,3,4-thiadiazole-2-sulfonamide as potential anti-pertussis drug using chromatography and spectroscopy techniques." *Transl Med (Sunnyvale)*, vol. 6, p. e138.
- [31] Heidari, A., 2016. "Nitrogen, oxygen, phosphorus and sulphur heterocyclic anti-cancer nano drugs separation in the supercritical fluid of ozone (o₃) using soave-redlich-kwong (srk) and pang-robinson (pr) equations." *Electronic J. Biol.*, vol. 12, p. 4.
- [32] Heidari, A., 2016. "An analytical and computational infrared spectroscopic review of vibrational modes in nucleic acids." *Austin J. Anal Pharm Chem.*, vol. 3, p. 1058.
- [33] Heidari, A., 2016. "Study of the role of anti-cancer molecules with different sizes for decreasing corresponding bulk tumor multiple organs or tissues." *Arch. Can. Res.*, vol. 4, p. 2.
- [34] Heidari, A., 2016. "Genomics and proteomics studies of zolpidem, necopidem, alpidem, saripidem, miroprofen, zolimidine, olprinone and abafungin as anti-tumor, peptide antibiotics, antiviral and central nervous system (cns) drugs." *J. Data Mining Genomics and Proteomics*, vol. 7, p. e125.
- [35] Heidari, A., 2016. "Pharmacogenomics and pharmacoproteomics studies of phosphodiesterase-5 (pde5) inhibitors and paclitaxel albumin-stabilized nanoparticles as sandwiched anti-cancer nano drugs between two DNA/rna molecules of human cancer cells." *J. Pharmacogenomics Pharmacoproteomics*, vol. 7, p. e153.
- [36] Heidari, A., 2016. "Biotranslational medical and biospectroscopic studies of cadmium oxide (CdO) nanoparticles-DNA/rna straight and cycle chain complexes as potent anti-viral, anti-tumor and anti-microbial drugs: A clinical approach." *Transl Biomed*, vol. 7, p. 2.
- [37] Heidari, A., 2016. "A comparative study on simultaneous determination and separation of adsorbed cadmium oxide (CdO) nanoparticles on DNA/rna of human cancer cells using biospectroscopic techniques and dielectrophoresis (dep) method." *Arch. Can. Res.*, vol. 4, p. 2.
- [38] Heidari, A., 2016. "Cheminformatics and system chemistry of cisplatin, carboplatin, nedaplatin, oxaliplatin, heptaplatin and lobaplatin as anti-cancer nano drugs: A combined computational and experimental study." *J. Inform Data Min.*, vol. 1, p. 3.
- [39] Heidari, A., 2016. "Linear and non-linear quantitative structure-anti-cancer-activity relationship (qsacar) study of hydrous ruthenium (iv) oxide (ruo₂) nanoparticles as non-nucleoside reverse transcriptase inhibitors (nnrtis) and anti-cancer nano drugs." *J. Integr Oncol*, vol. 5, p. e110.
- [40] Heidari, A., 2016. "Synthesis, characterization and biospectroscopic studies of cadmium oxide (CdO) nanoparticles-nucleic acids complexes absence of soluble polymer as a protective agent using nucleic acids condensation and solution reduction method." *J. Nanosci Curr. Res.*, vol. 1, p. e101.
- [41] Heidari, A., 2016. "Coplanarity and collinearity of 4'-dinonyl-2,2'-bithiazole in one domain of bleomycin and pingyangmycin to be responsible for binding of cadmium oxide (CdO) nanoparticles to DNA/rna bidentate ligands as anti-tumor nano drug." *Int. J. Drug Dev. and Res.*, vol. 8, pp. 007-008.
- [42] Heidari, A., 2016. "A pharmacovigilance study on linear and non-linear quantitative structure (chromatographic) retention relationships (qsrr) models for the prediction of retention time of anti-cancer nano drugs under synchrotron radiations." *J. Pharmacovigil*, vol. 4, p. e161.
- [43] Heidari, A., 2016. "Nanotechnology in preparation of semipermeable polymers." *J. Adv. Chem. Eng.*, vol. 6, p. 157.
- [44] Heidari, A., 2016. "A Gastrointestinal Study on Linear and Non-Linear Quantitative Structure (Chromatographic) Retention Relationships (QSRR) Models for Analysis 5-Aminosalicylates Nano Particles as Digestive System Nano Drugs under Synchrotron Radiations." *J. Gastrointest Dig. Syst.*, vol. 6, p. e119.
- [45] Heidari, A., 2016. "DNA/RNA fragmentation and cytolysis in human cancer cells treated with diphthamide nano particles derivatives." *Biomedical Data Mining*, vol. 5, p. e102.
- [46] Heidari, A., 2016 "A translational biomedical approach to structural arrangement of amino acids' complexes: A combined theoretical and computational study." *Transl Biomed*, vol. 7, p. 2.

- [47] Heidari, A. and Brown, C., 2015. "Study of composition and morphology of cadmium oxide (CdO) nanoparticles for eliminating cancer cells." *J. Nanomed Res.*, vol. 2, p. 20.
- [48] Heidari, A. and Brown, C., 2015. "Study of surface morphological, phytochemical and structural characteristics of rhodium (iii) oxide (rh₂o₃) nanoparticles." *International Journal of Pharmacology, Phytochemistry and Ethnomedicine*, vol. 1, pp. 15–19.
- [49] Heidari, A. and Brown, C., 2016. "Phase, composition and morphology study and analysis of os–pd/hfc nanocomposites." *Nano Res. Appl.*, vol. 2, p. 1.
- [50] Heidari, A. and Brown, C., 2016. "Vibrational spectroscopic study of intensities and shifts of symmetric vibration modes of ozone diluted by cumene." *International Journal of Advanced Chemistry*, vol. 4, pp. 5–9.